

Note. In this question-paper, \mathbb{R} denotes the set of real numbers.

1. Consider a board having 2 rows and n columns. Thus there are $2n$ cells in the board. Each cell is to be filled in by 0 or 1.
 - (a) In how many ways can this be done such that each row sum and each column sum is even?
 - (b) In how many ways can this be done such that each row sum and each column sum is odd?

2. Consider the function

$$f(x) = \sum_{k=1}^m (x - k)^4, \quad x \in \mathbb{R},$$

where $m > 1$ is an integer. Show that f has a unique minimum and find the point where the minimum is attained.

3. Consider the parabola $C : y^2 = 4x$ and the straight line $L : y = x + 2$. Let P be a variable point on L . Draw the two tangents from P to C and let Q_1 and Q_2 denote the two points of contact on C . Let Q be the mid-point of the line segment joining Q_1 and Q_2 . Find the locus of Q as P moves along L .
4. Let $P(x)$ be an odd degree polynomial in x with real coefficients. Show that the equation $P(P(x)) = 0$ has at least as many *distinct* real roots as the equation $P(x) = 0$.
5. For any positive integer n , and $i = 1, 2$, let $f_i(n)$ denote the number of divisors of n of the form $3k + i$ (including 1 and n). Define, for any positive integer n ,

$$f(n) = f_1(n) - f_2(n).$$

Find the values of $f(5^{2022})$ and $f(21^{2022})$.

6. Consider a sequence P_1, P_2, \dots of points in the plane such that P_1, P_2, P_3 are non-collinear and for every $n \geq 4$, P_n is the midpoint of the line segment joining P_{n-2} and P_{n-3} . Let L denote the line segment joining P_1 and P_5 . Prove the following:

- (a) The area of the triangle formed by the points P_n, P_{n-1}, P_{n-2} converges to zero as n goes to infinity.
- (b) The point P_9 lies on L .

7. Let

$$P(x) = 1 + 2x + 7x^2 + 13x^3, \quad x \in \mathbb{R}.$$

Calculate for all $x \in \mathbb{R}$,

$$\lim_{n \rightarrow \infty} \left(P\left(\frac{x}{n}\right) \right)^n.$$

8. Find the minimum value of

$$|\sin x + \cos x + \tan x + \cot x + \sec x + \operatorname{cosec} x|$$

for real numbers x not multiple of $\pi/2$.

9. Find the smallest positive real number k such that the following inequality holds

$$|z_1 + \dots + z_n| \geq \frac{1}{k} (|z_1| + \dots + |z_n|).$$

for every positive integer $n \geq 2$ and every choice z_1, \dots, z_n of complex numbers with non-negative real and imaginary parts.

[Hint: First find k that works for $n = 2$. Then show that the same k works for any $n \geq 2$.]