

**Indian Statistical Institute, Bangalore**  
M.S (QMS):2015

**Instructions**

The test is divided into two sessions (i) Forenoon session and (ii) Afternoon session. Each session is for two hours. For the forenoon session question paper, the test code is QMA and for the afternoon session question paper, the test code is QMB. Candidates appearing for MS(QMS) should verify and ensure that they are answering the right question paper.

The test QMA is multiple choice type. For each question, exactly one of the four choices is correct. You get four marks for each correct answer, one mark for each unanswered question, and zero mark for each incorrect answer.

The test QMB is of short answer type. It has altogether 10 questions. A candidate has to answer minimum of 8 questions.

Questions will be set on the following and related topics.

**Syllabus of QMA (Mathematics) and QMB (Mathematics)**

**Algebra:** Binomial Theorem, AP, GP, HP, Exponential, Logarithmic Series, Sequence, Permutations and Combinations, Theory of Equations.

**Matrix Algebra:** Vectors and Matrices, Matrix Operations, Determinants.

**Calculus:** Functions, Limits, Continuity, Differentiation of functions of one or more variables. Unconstrained Optimization, Definite and Indefinite Integrals: Integration by parts and integration by substitution. Elements of Probability and Probability distributions.

**Sample questions for QMA (Mathematics)(2015)**

1. If  $x = \frac{p}{(q-r)}$ ,  $y = \frac{q}{(r-p)}$ ,  $z = \frac{r}{(p-q)}$  with  $p \neq q \neq r$ , then the value of  $yz + zx + xy$  is
  - (a) 1
  - (b) 0
  - (c) -1
  - (d) 2
2. The value of  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2}$ , when  $u = \tan^{-1}(\frac{x}{y})$  is
  - (a) -1
  - (b) 0
  - (c) 1
  - (d) -2
3. The value of integral  $\int \int xy \, dx \, dy$ , over the region  $A$ , where  $A$  is the domain bounded by  $x$ -axis, ordinate  $x = 2a$  and the curve  $x^2 = 4ay$  is
  - (a)  $\frac{a^3}{2}$

- (b)  $\frac{a^4}{3}$
- (c)  $\frac{a^2}{4}$
- (d) 1

4.  $\int \frac{1}{x + x \log x} dx$  equals to

- (a)  $\log|x + \log x| + \text{Constant}$
- (b)  $\log|1 + x \log x| + \text{Constant}$
- (c)  $\log|\log x| + \text{Constant}$
- (d)  $\log|1 + \log x| + \text{Constant}$

5. The value of  $\frac{2}{3!} + \frac{4}{5!} + \frac{6}{7!} + \dots$  equals to

- (a)  $e$
- (b)  $\frac{1}{e^2}$
- (c)  $1 - \frac{1}{e}$
- (d)  $\frac{1}{e}$

6. The sales of a firm is promoted by advertisement campaign according to the rule  $\frac{ds}{dt} = 100e^{-0.25t}$  where  $S$  denotes the sales (in millions of rupees) and  $t$  is the number of years since the close of campaign. The sales promoted during the fifth year equals to

- (a) 30.56 millions of rupees
- (b) 31.44 millions of rupees
- (c) 32.32 millions of rupees
- (d) 32.56 millions of rupees

7. The Indian cricket team consists of 16 players. It includes 2 wicket keepers and 5 bowlers. In how many ways can a cricket team of eleven be selected if we have to select 1 wicket keeper and at least 4 bowlers?

- (a) 1200
- (b) 1056
- (c) 1072
- (d) 1092

8. Let  $\begin{vmatrix} a & a^2 & a^3 & -1 \\ b & b^2 & b^3 & -1 \\ c & c^2 & c^3 & -1 \end{vmatrix} = 0$ , in which  $a, b, c$  are all different. Then the value of  $abc$  is

- (a) 10
- (b) 1
- (c) 0.1
- (d) 0.01

9. The product of the first 100 positive integers ends with:

- (a) 21 zeroes
- (b) 22 zeroes
- (c) 23 zeroes
- (d) 24 zeroes

10. The function  $f(x) = |x| + \sin x + \cos^3 x$  is:
- continuous but not differentiable at  $x = 0$
  - differentiable at  $x = 0$
  - a bounded function which is not continuous at  $x = 0$
  - a bounded function which is continuous at  $x = 0$
11. Consider a cubical box of  $1m$  side which has one corner placed at  $(0,0,0)$  and the opposite corner placed at  $(1,1,1)$ . The least distance that an ant crawling from the point  $(0,0,0)$  to the point  $(1,1,1)$  must travel is:
- $\sqrt{6}m$
  - $\sqrt{5}m$
  - $2\sqrt{3}m$
  - $(1 + \sqrt{3})m$
12. The roots of the equation  $x^4 + x^2 = 1$  are
- all real and positive
  - never real
  - 2 positive and 2 negative
  - 1 positive, 1 negative and 2 non-real
13. I sold 2 books for Rs.30 each. My profit on one was 25 percent and the loss on the other was 25 percent. Then, on the whole, I
- lost Rs.5
  - lost Rs.4
  - gained Rs.5
  - neither gained nor lost
14. If  $a, b, c$  are real numbers so that  $x^3 + ax^2 + bx + c = (x^2 + 1)g(x)$  for some polynomial  $g$ , then
- $b = 1, a = c$
  - $b = 0 = c$
  - $a = 0$
  - none of the above
15. If  $E = \frac{1}{4} \cdot \frac{2}{6} \cdot \frac{3}{8} \cdot \frac{4}{10} \dots \frac{30}{62} \cdot \frac{31}{64} = 8^x$  then the value of  $x$  is
- 7
  - 9
  - 10
  - 12
16. If  $A$  and  $B$  are two square matrices such that  $B = -A^{-1}BA$ , then  $(A + B)^2$  is equal to
- 0
  - $A^2 + B^2$
  - $A^2 + 2AB + B^2$

(d)  $A + B$

17. In an examination of 9 papers, a candidate has to pass in more papers than the number of papers in which he fails in order to be successful. Find how many ways he can be successful.

(a) 256

(b) 216

(c) 220

(d) 272

18. If  $e^{e^x} = a_0 + a_1x + a_2x^2 + \dots$ , a power series, then

(a)  $a_0 = 1$

(b)  $a_0 = e$

(c)  $a_0 = e^e$

(d)  $a_0 = e^2$

19.  $\lim_{x \rightarrow 0} \frac{\sin(\pi \cos^2 x)}{x^2}$  equals to

(a)  $-\pi$

(b)  $\pi$

(c)  $\frac{\pi}{2}$

(d) 1

20. If there are three children in a family, then probability that there is one girl child in the family is

(a)  $\frac{2}{3}$

(b)  $\frac{1}{3}$

(c)  $\frac{1}{8}$

(d)  $\frac{3}{8}$

**Sample questions for QMB (Mathematics)(2015)**

1. (a) A set contains  $(2n + 1)$  elements. Find the number of subsets of the set which contain at most  $n$  elements.  
 (b) If  $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$  then show that  $A^n = \begin{bmatrix} \cos n\alpha & \sin n\alpha \\ -\sin n\alpha & \cos n\alpha \end{bmatrix}$
2. (a) Let  $f : R \rightarrow R$  such that for  $x, y \in R, |f(x) - f(y)| \leq (x - y)^2$ . Show that  $f$  is constant.  
 (b) If  $\alpha_1, \alpha_2, \dots, \alpha_n$  are the roots of the equation  $x^n + nax - b = 0$  then show that  $(\alpha_1 - \alpha_2)(\alpha_1 - \alpha_3)\dots(\alpha_1 - \alpha_n) = n(\alpha_1^{n-1} + a)$ .
3. (a) Suppose integers are formed by taking one or more digits from the following: 2,2,3,3,4,5,5,6,7. For example, 355 is a possible choice while 44 is not. Find the number of distinct integers that can be formed in which
  - i. the digits are non-decreasing;
  - ii. the digits are strictly increasing.
- (b) Show that 
$$\begin{vmatrix} x & l & m & 1 \\ \alpha & x & n & 1 \\ \alpha & \beta & x & 1 \\ \alpha & \beta & y & 1 \end{vmatrix} = (x - \alpha)(x - \beta)(x - y).$$
4. (a) Evaluate  $\int_0^1 \int_x^{\sqrt{2-x^2}} \frac{x}{\sqrt{x^2+y^2}} dy dx$ .  
 (b) Let  $A = ((a_{ij}))$  be an  $n \times n$  matrix such that  $a_{ij} = 0$  whenever  $i \geq j$ . Prove that  $A^n$  is a zero matrix.
5. (a) Two functions  $f$  and  $g$  are defined as follows for all non-negative real numbers.  $f(x) = \exp(-x)$  and  $g$  is the piece-wise linear function defined by connecting  $f(0), f(1), f(2), \dots$ . That is, for any non-negative integer  $k$ , the function  $g$  is obtained on the interval  $[k, k + 1)$  by connecting the points  $(k, f(k))$  and  $(k + 1, f(k + 1))$  with a line segment. Find the total area of the region which lies between the curves of  $f$  and  $g$ .  
 (b) For real numbers  $\alpha$  and  $\beta$ , consider the function  $f(x) = (\alpha - |x|)^2 e^{(\beta - |x|)^2}, -\infty < x < \infty$ .
  - i. For what values of  $\alpha$  and  $\beta$  is  $f$  continuous at all  $x$ ?
  - ii. For what values of  $\alpha$  and  $\beta$  is  $f$  differentiable at all  $x$ ?
6. (a) Find the value of  $\int_0^\infty \frac{\beta}{\eta} \left(\frac{x-\mu}{\eta}\right)^{\beta-1} \exp\left[-\left(\frac{x-\mu}{\eta}\right)^\beta\right] dx$  where  $\beta > 0, \eta > 0$ .  
 (b) Let  $g(x) = x^6 - x^5 + x^2 - x + 3, -\infty < x < \infty$ . Show that  $g(x) > 0$  for all  $x$ .
7. (a) Find the sum of the infinite series  $\left(\frac{1}{1.2} + \frac{1.3}{1.2.3.4} + \frac{1.3.5}{1.2.3.4.5.6} + \dots\right)$   
 (b) Show that  $1 < \int_0^1 e^{x^2} dx < e$
8. (a) Find two positive numbers such that their sum is equal to  $K$  and satisfy the property that the product of cube of the first number and square of the second number yields the maximum possible value.  
 (b) A cylindrical vessel of volume  $25\frac{1}{7}$  cubic metre, which is open at the top is to be manufactured from a sheet of metal. Find the dimensions of the vessel, so that the amount of sheet used in manufacturing it, is the least possible.
9. (a) If  $a = 11111\dots 1$  (63 digits),  $b = 1 + 10 + 10^2 + \dots + 10^6$  and  $c = 1 + 10^7 + 10^{14} + \dots + 10^{56}$ , then show that  $a = bc$ .

- (b) Suppose  $a, b$  and  $c$  are distinct non-zero real numbers and the roots of the equation  $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$  are equal, then show that  $a, b, c$  are in Harmonic Progression.
10. (a) Find the number of 5 digit numbers greater than 40000 that can be formed from the digits 0,1,2,3,4 and 5 when no repetitions are allowed.
- (b) For a positive integer  $n$ , express the function  $g(x) = (3 + x)^n$  as a polynomial:  $g(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$ . It is given that  $\sum_{j=0}^n a_j = 4096$ .
- Find the value of  $n$ .
  - Find the largest coefficient  $a_j$ .